

String Theory and Nonrelativistic Gravity

Eric Bergshoeff

Groningen University

work done in collaboration with

Ceyda Şimşek, Jaume Gomis, Kevin Grosvenor and Ziqi Yan

The 5th Conference of the Polish Society on General Relativity


Wojanów Palace, September 27, 2018



rijksuniversiteit
 groningen

Motivation

Holography



Gravity is not only used to describe the gravitational force!

Non-relativistic Holography

two approaches

- Keep general relativity in the bulk but take background geometry with **non-relativistic isometries**

Christensen, Hartong, Kiritsis, Obers and Rollier (2013-2015)

- Take **non-relativistic gravity** in the bulk

Gomis, Ooguri (2001); Gopakumar, Bagchi (2009)

Outline

String Newton-Cartan Gravity

Outline

String Newton-Cartan Gravity

The Non-relativistic Polyakov String

Outline

String Newton-Cartan Gravity

The Non-relativistic Polyakov String

T-duality

Outline

String Newton-Cartan Gravity

The Non-relativistic Polyakov String

T-duality

Actions

Outline

String Newton-Cartan Gravity

The Non-relativistic Polyakov String

T-duality

Actions

Comments

Outline

String Newton-Cartan Gravity

The Non-relativistic Polyakov String

T-duality

Actions

Comments

4D Galilei Symmetries

- time translations: $\delta t = \xi^0$ but not $\delta t = \lambda^i x^i$!
- space translations: $\delta x^i = \xi^i$ $i = 1, 2, 3$
- spatial rotations: $\delta x^i = \lambda^i_j x^j$
- Galilean boosts: $\delta x^i = \lambda^i t$

‘Conventional’ Constraints

symmetry	generators	gauge field	#	curvatures	#
time translations	H	τ_μ		$\tau_{\mu\nu} = \partial_{[\mu}\tau_{\nu]}$	6
space translations	P^A	E_μ^A		$R_{\mu\nu}^A(P)$	18
Galilean boosts	G^A	Ω_μ^A	12	$R_{\mu\nu}^A(G)$	
spatial rotations	J^{AB}	Ω_μ^{AB}	12	$R_{\mu\nu}^{AB}(J)$	

$$\mu = 0, 1, 2, 3; A = 1, 2, 3$$

$$R_{\mu\nu}^A(P) = 0 \quad (18) :$$

does only solve for **part of** $\Omega_\mu^A, \Omega_\mu^{AB}$

From Galilei to Bargmann

the **zero commutator**

$$[G_A, P_B] = 0$$

implies that a **massive particle** with non-zero spatial momentum P_B cannot by any boost transformation G_A be brought to a **rest frame** \Rightarrow

$$[G_A, P_B] = \delta_{AB} \mathbf{Z} \quad \rightarrow \quad \text{extra gauge field } m_\mu$$

and additional **6** conventional constraints: $R_{\mu\nu}(Z) = 0$

'Geometric' Constraints

$$\tau_{\mu\nu} \equiv \partial_{[\mu} \tau_{\nu]} = 0 \quad \rightarrow \quad \tau_{\mu} = \partial_{\mu} \rho \quad \text{with} \quad \tau_{\mu} \text{ clock function}$$



$$\Delta T = \int_C dx^{\mu} \tau_{\mu} = \int_C d\rho \text{ is path-independent} \quad \rightarrow \quad \text{absolute time}$$

NC Gravity couples to **particles**

what about **strings**?

Comparing Limits

The NR limit of a particle coupled to general relativity is **finite** provided that the particle is coupled to a **(central charge) gauge field B_μ** which is expressed in terms of the **timelike NC Vierbein field τ_μ** as

$$B_\mu = \omega \tau_\mu$$

with $\omega \rightarrow \infty$ in the NR limit

Similarly, the string should be coupled to a **2-form gauge field $B_{\mu\nu}$** with

$$B_{\mu\nu} = \omega^2 \epsilon_{AB} \tau_\mu^A \tau_\nu^B, \quad A = 0, 1$$

defining a **string NC geometry** with **generalized clock functions τ_μ^A**

see also Gomis, Ooguri (2001)

String Galilei Symmetries

Gomis, Ooguri (2001); Andringa, Gomis, de Roo + E.B. (2012)

$$D + 1 \text{ flat indices} \rightarrow \begin{cases} 2 \text{ longitudinal indices } A \\ D - 1 \text{ transverse indices } A' \end{cases}$$

longitudinal translations H_A

transverse translations $P_{A'}$

string Galilei boosts $G_{AB'}$

longitudinal Lorentz rotations M_{AB}

transverse spatial rotations $J_{A'B'}$

Conventional versus Geometric Constraints

$$R_{\mu\nu}{}^A(H) = D_{[\mu}(\Omega)\tau_{\nu]}{}^A = 0 \quad \text{with} \quad \tau_{\mu}{}^A \text{ generalized clock function}$$

4D : 12 components

- 4 conventional constraints: solves for $\Omega_{\mu}{}^{AB} = \epsilon^{AB}\Sigma_{\mu}$
- 8 geometric constraints

Noncentral Extension

$$[G_{AA'}, P_{B'}] = 0 \quad \rightarrow \quad [G_{\textcolor{red}{A}A'}, P_{B'}] = \delta_{A'B'} \textcolor{red}{Z}_A$$

The independent string NC fields $\{\tau_\mu^A, e_\mu^{A'}, m_\mu^A\}$ transform as follows:

$$\begin{aligned} \delta \tau_\mu^A &= \Lambda^A_B \tau_\mu^B, \\ \delta E_\mu^{A'} &= \Lambda^{A'}_{B'} E_\mu^{B'} - \textcolor{red}{\Sigma}_A^{A'} \tau_\mu^A, \\ \delta m_\mu^A &= D_\mu \sigma^A + \textcolor{red}{\Sigma}_{A'}^A E_\mu^{A'} \end{aligned}$$

longitudinal metric:

$$\tau_{\mu\nu} \equiv \tau_\mu^A \tau_\nu^B \eta_{AB}$$

transverse ‘metric’:

$$H_{\mu\nu} \equiv E_\mu^{A'} E_\nu^{B'} \delta_{A'B'} + (\tau_\mu^A m_\nu^B + \tau_\nu^A m_\mu^B) \eta_{AB}$$

Outline

String Newton-Cartan Gravity

The Non-relativistic Polyakov String

T-duality

Actions

Comments

NR Limit Polyakov Particle

$$S_{\text{Pol.}} = -\frac{1}{2} \int d\tau \left\{ -\frac{1}{e} E_{\mu}^A \dot{x}^{\mu} E_{\nu}^B \dot{x}^{\nu} \eta_{AB} + M^2 e - 2M B_{\mu} \dot{x}^{\mu} \right\}$$

$$\begin{aligned} S_{\text{Pol.}}(c^2) &= -\frac{1}{2} \int d\tau \frac{1}{e} c^2 [\tau_{\mu} \dot{x}^{\mu} - me]^2 \\ &= -\frac{1}{2} \int d\tau \frac{1}{e} \left\{ \lambda (\tau_{\mu} \dot{x}^{\mu} - me) - \frac{1}{4c^2} \lambda^2 \right\} \Rightarrow \end{aligned}$$

$$S_{\text{Pol.}}(\text{N.R.}) = -\frac{1}{2} \int d\tau \frac{1}{e} \left\{ \dot{x}^{\mu} \dot{x}^{\nu} H_{\mu\nu} + \lambda (\tau_{\mu} \dot{x}^{\mu} - me) \right\}$$

Nambu-Goto (NG)

- **relativistic NG** is non-linear in **longitudinal** and **transverse** embedding coordinates
- **nonrelativistic NG** is nonlinear in **longitudinal** embedding coordinates only

The Nonrelativistic Polyakov String

$$h_{\alpha\beta} = e_{\alpha}{}^a e_{\beta}{}^b \eta_{ab}$$

$$e_{\alpha} \equiv e_{\alpha}{}^0 + e_{\alpha}{}^1, \quad \bar{e}_{\alpha} \equiv e_{\alpha}{}^0 - e_{\alpha}{}^1$$

$$\tau_{\mu} \equiv \tau_{\mu}{}^0 + \tau_{\mu}{}^1, \quad \bar{\tau}_{\mu} \equiv \tau_{\mu}{}^0 - \tau_{\mu}{}^1$$

$$S_{\text{Pol.}} = -\frac{T}{2} \int d^2\sigma \left[\sqrt{-h} h^{\alpha\beta} \partial_{\alpha} x^{\mu} \partial_{\beta} x^{\nu} H_{\mu\nu} + \epsilon^{\alpha\beta} (\lambda e_{\alpha} \tau_{\mu} + \bar{\lambda} \bar{e}_{\alpha} \bar{\tau}_{\mu}) \partial_{\beta} x^{\mu} \right]$$

$$- \frac{T}{2} \int d^2\sigma \epsilon^{\alpha\beta} \partial_{\alpha} x^{\mu} \partial_{\beta} x^{\nu} B_{\mu\nu}$$

Nambu-Goto : $e_{\alpha}{}^a \neq \tau_{\alpha}{}^A$ but $h_{\alpha\beta} = \tau_{\alpha\beta}$ up to a scale factor

World-sheet Analysis

Gomis, Ooguri (2001)

Conformal Gauge

$$\sqrt{-h} h^{\alpha\beta} = \eta^{\alpha\beta}$$

Flat Spacetime

$$\tau_{\mu}{}^A = \delta_{\mu}^A, \quad E_{\mu}{}^{A'} = \delta_{\mu}^{A'}, \quad m_{\mu}{}^A = 0$$

\Downarrow

$$S = -\frac{T}{2} \int d^2\sigma \left(\partial X^{A'} \bar{\partial} X^{B'} \delta_{A'B'} + \lambda \bar{\partial} X + \bar{\lambda} \partial \bar{X} \right)$$

Outline

String Newton-Cartan Gravity

The Non-relativistic Polyakov String

T-duality

Actions

Comments

Relativistic T-duality

Buscher (1987,1988); Roček, Verlinde (1992)

adapted coordinates: $x^\mu = (y, x^i)$ $k^\mu \partial_\mu = \partial_y$

$$S_{\text{parent}} = \underbrace{S_{\text{Pol.}}(\partial_\alpha y \rightarrow v_\alpha)}_{\text{quadratic in } v_\alpha!} - T \int d^2 \sigma \epsilon^{\alpha\beta} \tilde{y} \partial_\alpha v_\beta$$

$$\frac{\delta S_{\text{parent}}}{\delta v_\alpha} = 0 \quad \rightarrow \quad v_\alpha \text{ is solved for in terms of dual coordinate } \tilde{y}$$

$$\tilde{G}_{yy} = \frac{1}{G_{yy}} \quad \Rightarrow \quad R \Leftrightarrow \frac{1}{R}$$

Non-relativistic Longitudinal T-duality

$$\tau_{\mu}^0 k^{\mu} = 0, \quad \tau_{\mu}^1 k^{\mu} \neq 0, \quad E_{\mu}^{A'} k^{\mu} = 0$$

adapted coordinates: $x^{\mu} = (y, x^i) \quad k^{\mu} \partial_{\mu} = \partial_y$

$$S_{\text{parent}} = S_{\text{Pol.}}(\partial_{\alpha} y \rightarrow v_{\alpha}) - T \int d^2 \sigma \epsilon^{\alpha\beta} \tilde{y} \partial_{\alpha} v_{\beta}$$

$$\frac{\delta S_{\text{parent}}}{\delta v_{\alpha}} = 0 \quad \rightarrow \quad v_{\alpha} \text{ is solved for in terms of dual coordinate } \tilde{y} \text{ and}$$

the Lagrange multipliers $\lambda, \bar{\lambda}$ if we dualize in the longitudinal direction

$\rightarrow \lambda, \bar{\lambda}$ are no Lagrange multipliers anymore!

Dual Action

$$\tilde{S}_{\text{long.}} = -\frac{T}{2} \int d^2\sigma \left(\sqrt{-h} h^{\alpha\beta} \partial_\alpha \tilde{x}^\mu \partial_\beta \tilde{x}^\nu \tilde{G}_{\mu\nu} + \epsilon^{\alpha\beta} \partial_\alpha \tilde{x}^\mu \partial_\beta \tilde{x}^\nu \tilde{B}_{\mu\nu} \right)$$

with $\tilde{x}^\mu = (\tilde{y}, x^i)$ and $\tilde{G}_{yy} = 0$: **lightlike direction**

- The **longitudinal spatial** T-dual of the NR string is the Polyakov string moving in a GR background with a **lightlike direction**
 - **DLCQ relativistic string**
 - **Rigid susy in Lorentzian curved spacetimes**

See, e.g., Cassani, Klare, Martelli, Tomasiello, Zaffaroni (2014)

- The **transverse spatial** T-dual of the NR string is again a NR string with a transverse spatial isometry direction à la Buscher

Outline

String Newton-Cartan Gravity

The Non-relativistic Polyakov String

T-duality

Actions

Comments

3D Extended Bargmann Gravity

$$S = \frac{1}{2\kappa^2} \int d^3x (E R(\Omega) - \epsilon^{\mu\nu\rho} B_\mu \partial_\nu A_\rho) \Rightarrow$$

Chern-Simons action for **3D Extended Bargmann Algebra**

Rosseel + E.B. (2016); Hartong, Obers (2016)

- The **2 central charges** originate from the **2 vector fields**
- **3D Extended Bargmann Gravity** \neq **3D NC gravity**

4D Extended String Bargmann Gravity

$$S = \frac{1}{2\kappa^2} \int d^4x (E R(\Omega) - \epsilon^{\mu\nu\rho\sigma} B_{\mu\nu} \partial_\rho A_\sigma)$$

Grosvenor, Şimşek, Yan + E.B., work in progress

- The 2-form gives rise to the **2 noncentral extensions** Z_A ($A = 0, 1$) of the String Bargmann Algebra
- The 1-form gives rise to the **extra central extension** S of the **Extended String Bargmann Algebra**

A 4D First-Order Action

Grosvenor, Şimşek, Yan + E.B. (to appear)

$$S = \frac{1}{2\kappa^2} \int d^4x \epsilon^{\mu\nu\rho\sigma} \left[-\frac{1}{2} \epsilon_{A'B'} E_\mu^{A'} E_\nu^{B'} \mathcal{R}_{\rho\sigma}(M) - \epsilon_{AB} \epsilon_{A'B'} \tau_\mu^A E_\nu^{A'} \mathcal{R}_{\rho\sigma}{}^{BB'}(G) \right. \\ \left. + \epsilon_{AB} \tau_\mu^A m_\nu^B \mathcal{R}_{\rho\sigma}(J) + \frac{1}{2} \epsilon_{AB} \tau_\mu^A \tau_\nu^B \mathcal{R}_{\rho\sigma}(S) \right]$$

Relation to 3D Extended Bargmann Gravity

The 3D Chern-Simons action of **Extended Bargmann gravity** is obtained after a dimensional reduction over the **longitudinal** y -direction with $\mu = (y, i)$, $i = 0, 1, 2$ followed by a **truncation**. For instance:

$$\tau_{\mu}^A = \begin{pmatrix} \tau_y^0 & \tau_y^1 \\ \tau_i^0 & \tau_i^1 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ \tau_i & 0 \end{pmatrix} \quad \Rightarrow$$

$$S_{\text{CS}} = \frac{k}{4\pi} \int d^3x \epsilon^{ijk} \left[\epsilon_{A'B'} E_i^{A'} \mathcal{R}_{jk}{}^{B'}(G) - m_i \mathcal{R}_{jk}(J) - \tau_i \mathcal{R}_{jk}(S) \right]$$

Outline

String Newton-Cartan Gravity

The Non-relativistic Polyakov String

T-duality

Actions

Comments

Nonrelativistic String Theory

- **Nonrelativistic String Theory** can be defined independent of any limit of **relativistic string theory**
 - T-duality
 - Target Space Actions
- **String NC Geometry** is to NR string theory what **Riemannian geometry** is to relativistic string theory

Generalization to $p + 3$ dimensions

In $p + 3$ dimensions the divergencies originating from the Einstein-Hilbert term are cancelled by introducing a $p + 1$ -form gauge field B and a 1-form gauge field A :

$$\epsilon^{\mu_1 \cdots \mu_{p+3}} \mathcal{B}_{\mu_1 \cdots \mu_{p+1}} \partial_{\mu_{p+2}} \mathcal{A}_{\mu_{p+3}}$$

The non-relativistic limit leads to an action for **extended p -brane Bargmann gravity** or **extended co-dimension 2 Bargmann gravity**

They are all related by dimensional reduction of a spatial direction of the p -brane followed by a truncation.

Open Issues

- Does β -function calculation leads to consistent backgrounds?

- Nonrelativistic holography?

Gopakumar, Bagchi (2009)

- Anyonic strings?

cp. to Duval, Horvathy (2000); Jackiw, Nair (2000); Mezincescu, Townsend (2010)

See also, Gußmann, Sarkar and Wintergerst (2018)

- Lie algebra expansions

Izquierdo, private communication

- Double Field Theory?

S. M. Ko, C. Melby-Thompson, R. Meyer and J.-H. Park (2015)

Take Home Message

Nonrelativistic String Theory can be studied!

Gomis, Ooguri (2001)